Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Microbiology Research

Journal homepage: <https://www.ijmronline.org/>

Original Research Article

Spectrum of surgical site infections at a tertiary care hospital in Hyderabad

Ahmed Abdul Muqtadir¹, Guruprasad Mandevwad¹, Rajkumar H R V¹,
Ruturaj M K¹, Sandeep Kumar Tipparthi¹, Ravi Shankar Reddy^{1,*}¹Dept. of Microbiology, Kamineni Academy of Medical Sciences and Research Centre, Bahadurguda, Telangana, India

ARTICLE INFO

Article history:

Received 27-09-2020

Accepted 16-11-2020

Available online 25-12-2020

Keywords:

Bacterial drug resistance

Surgical site infections

ABSTRACT

Introduction: Surgical site infections (SSI) constitute a major public health problem worldwide and are the second most frequently reported nosocomial infections. They are responsible for increasing the treatment cost, readmission, length of hospital stay and significant morbidity and mortality. The present study was aimed to determine the incidence of SSIs and the prevalence of aerobic bacterial pathogens involved with their antibiogram at our tertiary care hospital.

Materials and Methods: This study was conducted for a period of one year (January to December 2019). The SSI occurrence, the organisms isolated, antibiotic sensitivity pattern as well as outcome of the treatment is evaluated.

Results: In overall surgeries conducted in a year at our institute, we found 15 cases from various department were clinically diagnosed of having SSIs, including 11 male and 4 females with the mean age of 43.7 years. Out of the total 15 samples processed, 9 (60%) yielded bacterial growth including *Staphylococcus species*, *Enterococcus*, *Pseudomonas* and *Acinetobacter species*. Most of these are organisms isolated were not multidrug resistant.

Discussion: Management of SSIs remains a significant concern for surgeons and physicians in a health care facility, which carries a load with high morbidity and mortality. Due to the proper hospital infection control management we report the occurrence of low SSI in our hospital with the absence of multidrug resistance. We emphasize the importance of hospital infection control monitoring with proper precautions during surgeries to reduce the load of SSI and better outcome of the treatment.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Surgical Site infections (SSI) is considered as a significant problem in clinical aspects associated with the high morbidity and mortality. It has been associated with greater economic burden, prolonged hospitalization and high mortality rate.¹ CDC clearly defined SSI as an infection that occurs after surgery in the part of the body where the surgery took place and classified the SSI into superficial, deep and organ specific.¹ The wide range of incidence and prevalence of SSI has been noted in United States and it is the most common complication occurring in

2-5% of post-surgery cases.² The pathogenesis of SSI has been associated both with the endogenous contamination such as skin flora and exogenous contamination by healthcare personnel or contaminated surgical instruments. Other factors such as burden of organism and virulence factors of pathogens also play a major role in occurrence of SSI.³ The risk factors associated with the SSI were classified as intrinsic associated with age, obesity, nutrition, preoperative hospitalization. The perioperative factors such as preparation of patient, hair removal, skin preparation, antimicrobial prophylaxis and duration of surgery impact the outcome of surgery and ill preparation favours the SSI.³

Literature revealed several microorganisms are associated with SSI and etiological pathogen has been

* Corresponding author.

E-mail address: dravishankarreddy@gmail.com (R. S. Reddy).

associated with the type of surgery. Owen et al. found that the common organisms associated with SSI are mostly gram positive bacteria such as *Staphylococcus aureus*, *coagulase negative Staphylococcus*, *Enterococcus species*.⁴ Survey of literature found that Indian studies have also contributed and found different pathogenic microorganisms associated with SSI. A study from Raj Kumari and co-workers found 4.4% of orthopaedic cases were confirmed as SSI and etiological agents included predominantly *Acinetobacter baumannii* and *Staphylococcus aureus*.⁵ A study from Uttarakhand has found the predominantly SSI has been associated with *Staphylococcus aureus* at the rural healthcare unit.⁶ Patak et al. in their detail study found statistically significant risk factors such as severity of the disease, presence of drains, history of previous hospitalization, preoperative stay and surgical duration associated with SSI.⁷ A study from Mumbai pointed out the serious concern of presence of multidrug resistant gram negative bacteria associated with SSI.⁸ To best of our knowledge, not many studies have been done in the Southern part of India to evaluate the occurrence of SSI and the microorganisms associated with it. The aim of the study is to evaluate the occurrence of SSI, associated comorbidities, the etiological microorganisms and their drug sensitivity pattern at our tertiary care centre.

2. Materials and Methods

This prospective study was done in the Department of Microbiology, Kamineni Academy of Medical Sciences and Research Centre, Hyderabad. The study period was from January 2019 to December 2019. The patient details were recorded including type of surgery, type of wound infection, wound class including clean, clean contaminated and contaminated wound and total days of stay in the hospital.(Table 1) (Figure 1)

2.1. Inclusion criteria

All the patients admitted from different surgical wards of the hospitals during the tenure of study period for elective or emergency surgeries associated with clean, clean contaminated and contaminated were included.

The demographic data, associated comorbidities, risk factors, duration of surgery and clinical evaluation of wound were collected and analysed.

2.2. Specimen collection

Cases categorised as SSI, specimens were collected from different surgical infected sites as mentioned in table 1. Pre-existing morbidities such as hypertension, chronic kidney disease (CKD), coronary artery disease (CAD) and various signs of SSI were noted.

Fig. 1: Exploratory laparotomy

2.3. Microbiological evaluation

All the samples were cultured on blood agar, MacConkey Agar media and Chromogenic agar media, and culture plates were incubated at 37°C overnight, the identification of culture and sensitivity to various antibiotics was carried out using VITEK-2 system.

Statistical analysis is done using Microsoft excel.

3. Results

In this period in our hospital a total number of 1613 surgeries were carried out, in which orthopaedic surgeries accounted for 632, gynaecology department accounted for 436 surgeries, surgery department 416 and 129 surgeries were carried out in Neurosurgery department. The patients were examined preoperatively, intra-operatively and post operatively. Out of 1613 cases the present study found the occurrence of 15 cases (0.92%) developed SSI (Figure 2). The patients included 11 males 4 females, the age of the patients were in a range of 17 years to 70 years. 5 patients were in age group 17-34, 5 patients in age group of 35-51, and 5 patients in an age group of 52-70 years with mean age of 43.7 years. The duration of the surgery lasting less than 2 hours has been noticed in 13 cases and in remaining two cases the duration of surgery was more than 2 hours. Few cases are with the comorbidities such as 3 hypertension cases, 1 case of chronic kidney disease, 1 case of coronary artery disease and 1 case has been admitted with road traffic accident. The cases taken up for the surgery were included as elective in 10 cases, and 5 cases were taken up on emergency basis (Figure 3).

Table 1: Details of type of surgery, type of wound, hospital stay and type of specimen collected

S.No	Surgery	Type of wound infections	Stay at Hospital (days)	Specimen
1	Exploratory laparotomy	Clean contaminated	5	Wound swab
2	Lower segment caesarean section	Clean contaminated	8	Wound swab
3	Lower segment caesarean section	Clean contaminated	8	Wound swab
4	Total abdominal Hysterectomy	Clean contaminated	11	Wound swab
5	Total abdominal Hysterectomy	Contaminated	48	Wound swab
6	Exploratory laparotomy	Clean contaminated	11	Wound swab
7	Open reduction internal fixation	Clean contaminated	8	Wound swab
8	Closed reduction internal fixation	Clean contaminated	11	Wound swab
9	Open reduction internal fixation	Clean contaminated	8	Tissue
10	Open reduction internal fixation	Clean contaminated	9	Pus
11	Open reduction internal fixation	Clean contaminated	6	Tissue
12	Minimal invasive percutaneous plate osteosynthesis	Clean contaminated	7	Pus
13	Anterior cruciate ligament reconstruction	Clean contaminated	5	Wound swab
14	Decompressive craniotomy site	Clean contaminated	9	Wound swab
15	Laminectomy	Clean Wound	50	Pus

Fig. 2: Represent incidence of SSI **Fig. 3:** Graph represents factors associated with SSI

These 15 cases presented with the signs and symptoms of surgical wound infections. Out of 15 cases a majority of 13 cases were clean contaminated wounds and 1 was clean wound and 1 was reported as contaminated wound. A total of 9 cases (60%) of SSI were culture positive out of 15 cases. From the elective procedures 4 cases (40%) were culture positive and in emergency procedures of 5 cases all were culture positive (100%). Among comorbid patients with hypertension, CKD, CAD all cultures were positive for

the bacterial growth.

The organisms isolated were *Staphylococcus species* which includes 3 MSSA (*Methicillin sensitive Staphylococcus aureus*), 1 MRSA (*methicillin resistant Staphylococcus aureus*) and 1 *Staphylococcus hemolyticus*, two *Enterococcus* species which includes one *Enterococcus durans* and another *Enterococcus faecalis*. Among gram negative bacteria that were isolated which are specific to SSI included *Pseudomonas aeruginosa* and the other showed

growth of *Acinetobacter baumannii* (Figure 4). Most of the organisms isolated were sensitive to several antibiotics except *Acinetobacter baumannii* which was resistant to several antibiotics (Figure 5). *Enterococcus species* and methicillin sensitive *Staphylococcus aureus* were sensitive to erythromycin, gentamicin, ciprofloxacin, linezolid, clindamycin and vancomycin. MRSA was sensitive to vancomycin, teicoplanin, amikacin, linezolid. In gram negative bacteria *Pseudomonas aeruginosa* was sensitive to piperacillin/tazobactam, amikacin, colistin, ceftazidime, imipenem, meropenem. *Acinetobacter baumannii* was found to be sensitive to colistin and tigecycline only.

Fig. 4: The spectrum of bacterial isolated from SSI cases

Fig. 5: A: *Staphylococcus aureus* on blood agar, B: *Pseudomonas aeruginosa* on CPSE agar

4. Discussion

Several studies all over the world has reported surgical site infection as one of the causative agent for the long stay in hospital, leading to spread of bacterial drug resistance as well as the psychological and monetary effect on the patients. The overall infection rate in the present study was less than 1%, a very less occurrence of SSI compared to the other studies which reported rates ranging from 2.5 to 41.9%.^{9,10} In the present study we report the occurrence of SSI at very low percentage which emphasize the continuity of strict adherence of standard operating procedure and

effective work of hospital infection control committee. In the present study the overall average stay in the hospital on average was 13.6 days (min-5 days- max-50 days). Similar to other studies the present study also found the most common bacterial isolates were *Staphylococcus species* with varying antibiotic sensitivity pattern.² Contrast to a study conducted by Lilani et al., which found occurrence of more gram negative bacteria isolated from clean-contaminated wounds, the present study isolated a single *Pseudomonas aeruginosa* and a single *Acinetobacter baumannii* complex.¹¹

The present study also emphasize the importance of duration of surgery. Emergency surgery is one of the key factors, and have more than 50% risk of developing SSI compared to elective procedures,¹² the present study also support and shown the 100% bacterial growth in SSI cases taken up on emergency basis. Comorbidities including hypertension, CKD and CAD also play a major and independent risk factors in SSI. A study from Mejía et al. has found the patients with comorbidities acts as risk factors favouring SSI and our results echo with similar results where we found 100% growth in SSI in cases having comorbidities as a one of the important factor.¹³ Age is one of the factors, which might increase the chances of having SSI. A cohort study conducted by Kaye et al has found that age is an independent factor which favours the occurrence of SSI, similar to our results which found high occurrence of culture positive cases in patients with age ranging from 52-70 compared to lower age group.¹⁴

The present study has found low SSI cases along with the organisms which were sensitive to most of the antibiotics except for *Acinetobacter baumannii* complex. The organisms were found sensitive to the several classes of antibiotics such as macrolides, cephalosporin's, beta lactamases, aminoglycosides, glycopeptides and carbapenems. This study also emphasise the importance of isolation of newer organisms in relation with the SSI. We first time report to our knowledge- the isolation of *Staphylococcus hemolyticus*, *Enterococcus durans* from the surgical site infection which are sensitive to most of the antibiotics. We isolated multidrug resistant *Acinetobacter baumannii* which has been associated with long stay in the hospital. We conclude the importance of surveillance of surgical site infection, proper pre and postsurgical management, strict adherence to hospital infection control guidelines to prevent the occurrence of SSI in tertiary care hospitals as well as prevention of bacterial drug resistance.

5. Source of Funding

None.

6. Conflict of Interest

The author(s) declare(s) that there is no conflict of interest regarding the publication of this article.

References

1. Kolasinski W. Surgical site infections - review of current knowledge, methods of prevention. *Pol Przegl Chir.* 2019;91(4):41–7.
2. Prabhakar H, Arora S. A bacteriological study of wound infections. *J Indian Med Assoc.* 1979;73:145–8.
3. Garner BH, Anderson DJ. Surgical Site Infections. *Infect Dis Clin N Am.* 2016;30(4):909–29. doi:10.1016/j.idc.2016.07.010.
4. Owens CD, Stoessel K. Surgical site infections: epidemiology, microbiology and prevention. *J Hosp Infect.* 2008;70(2):3–10. doi:10.1016/s0195-6701(08)60017-1.
5. Mathur P, Trikha V, Sharma V, Farooque K, Misra MC, Rajkumari N, et al. Outcomes of surgical site infections in orthopedic trauma surgeries in a tertiary care centre in India. *J Postgrad Med.* 2014;60(3):254. doi:10.4103/0022-3859.138731.
6. Juyal D, Pal S, Sayana A, Joshi A. Staphylococcus aureus: A predominant cause of surgical site infections in a rural healthcare setup of Uttarakhand. *J Fam Med Prim Care.* 2019;8(11):3600. doi:10.4103/jfmprc.jfmprc_521_19.
7. Pathak A, Saliba EA, Sharma S, Mahadik VK, Shah H, Lundborg CS. Incidence and factors associated with surgical site infections in a teaching hospital in Ujjain, India. *Am J Infect Control.* 2014;42(1):e11–5. doi:10.1016/j.ajic.2013.06.013.
8. Shah S, Singhal T, Naik R. A 4-year prospective study to determine the incidence and microbial etiology of surgical site infections at a private tertiary care hospital in Mumbai, India. *Am J Infect Control.* 2015;43(1):59–62. doi:10.1016/j.ajic.2014.10.002.
9. Agarwal SL. Study of postoperative wound infection. *Indian J Surg.* 1972;34:314–20.
10. Anvikar AR, Deshmukh AB, Karyakarte RP, Damle AS, Patwardhan NS, Malik AK. A one year prospective study of 3,280 surgical wounds. *Indian J Med Microbiol.* 1999;17:129–32.
11. Lilani SP, Jangale N, Chowdhary A, Daver GB. Surgical site infection in clean and clean-contaminated cases. *Indian J Med Microbiol.* 2005;2(4):249–52.
12. Rubin RH. Surgical wound infection: epidemiology, pathogenesis, diagnosis and management. *BMC infect Dis.* 2006;6:171.
13. Mejía MPI, Verduzco JMF, López MLL, Acosta MEH, Silva JD. Patient comorbidities as risk factors for surgical site infection in gynecologic and obstetric surgery. *Int J Fam Community Med.* 2019;3(2):91–4. doi:10.15406/ijfcm.2019.03.00137.
14. Kaye KS, Schmit K, Pieper C, Sloane R, Caughlan KF, Sexton DJ, et al. The Effect of Increasing Age on the Risk of Surgical Site Infection. *J Infect Dis.* 2005;191(7):1056–62. doi:10.1086/428626.

Author biography

Ahmed Abdul Muqtadir, Post Graduate Student

Guruprasad Mandevwad, Assistant Professor

Rajkumar H R V, Professor and HOD

Ruturaj M K, Assistant Professor

Sandeep Kumar Tippiarthy, Tutor

Ravi Shankar Reddy, Professor

Cite this article: Muqtadir AA, Mandevwad G, Rajkumar H R V, Ruturaj M K, Tippiarthy SK, Reddy RS. Spectrum of surgical site infections at a tertiary care hospital in Hyderabad. *Indian J Microbiol Res* 2020;7(4):322-326.