Original Research Article

Detection of catheter related blood stream infections in an ICU of a tertiary care center in Northern India

Humaira Basher¹, Anjum Farhana^{1,*}, Seema Alim¹, Shoiab Khan¹¹Dept. of Microbiology, Government Medical College, Srinagar, Jammu & Kashmir, India

ARTICLE INFO

Article history:

Received 12-05-2021

Accepted 17-05-2021

Available online 30-07-2021

Keywords:

Microorganisms

Bacterial biofilms

ABSTRACT

Introduction: Indwelling medical devices are frequently used in all health setting with critical care units of hospitals for treatment and intervention in patient care. Microorganisms attach to surfaces resulting in the formation of a biofilm which pose a serious public health issue because of its increased resistance to antimicrobial agents and the potential to cause infections.

Aims and Objectives: To determine the proportion of bacterial Biofilms in patients with central venous line and to find out the organisms most commonly associated with it.

Materials and Methods: The study was carried out from January 2019 to September 2020. Positive cultures were obtained from 58 of these 102 samples. The isolates were then tested for in vitro production of biofilm using a microtitre plate assay. All the isolates were subjected to antimicrobial susceptibility tests on Muller-Hinton agar by Kirby Bauer disc diffusion method based on CLSI guidelines.

Results: Demographic characteristics of the study subjects showed that among 102 catheterized patients isolated with males 58.82% and 41.1% females. Out of 58 isolates, 62.06% were gram negative bacteria (GNB) and 37.93% were gram positive. *Staphylococcus aureus* was most common isolate 27.58% followed by *Klebsiella spp.* 20.68%, *Acinetobacter* 15.5%, *Coagulase negative Staphylococci* 6.8% respectively. The total number of positive slime producers, in this study, was 23(39.6%). The highest number of strong slime producer strains was observed in case of *Pseudomonas* (2 out of 5), *Klebsiella pneumoniae* (4 out of 12) and *Staphylococcus* (3 out of 16). Antibiotic susceptibility patterns showed increased resistance towards penicillin and beta -lactam group of antibiotics, increased sensitivity to linezolid and vancomycin among gram positive organisms. Among gram negative bacteria increased resistance was seen for cephalosporins and aminoglycosides and least resistance for colistin.

Conclusion: Colonization of indwelling medical devices with consequent biofilm production is a likely contributory factor to infections. The microorganisms survive in the hospital environment despite unfavourable conditions such as desiccation, nutrient starvation, and antimicrobial treatment. Resistance to antibiotics ladder is increasing and it's necessary to take actions to reduce its hindrance in the future. Advanced studies in biofilm will help to prevent the more virulent factors which protect the bacteria from host immunity and to get rid of critical complications in therapy.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Microorganisms produce extracellular polysaccharides, resulting in the formation of a biofilm that attach to surfaces. Because of the increased resistance of biofilm-associated organisms to antimicrobial agents they pose

a serious problem for public health. There is a potential for these organisms to cause infections in patients with indwelling medical devices. Many bloodstream infections and urinary tract infections are associated with indwelling medical devices and, therefore, are (in most cases) biofilm associated. The most effective strategy for treating these infections may be removal of the biofilm contaminated device.¹

* Corresponding author.

E-mail address: anjumfarhana1@gmail.com (A. Farhana).

A biofilm is defined as an architectural colony of microorganisms, within a matrix of extracellular polymeric substance that they produce. Biofilm contains microbial cells adherent to one-another and to a static surface (living or non-living). Biofilms are usually pathogenic in nature and are associated with nosocomial infections. The National Institutes of Health (NIH) has shown that 65% of microbial and 80% of chronic infections are associated with biofilm formation. There are many steps in biofilm formation, starting with attachment to a living or non-living surface that will lead to formation of micro-colony, giving rise to three-dimensional structures and ending up, after maturation, with detachment. Several species of bacteria communicate with one another during biofilm formation, through quorum sensing. In general, bacterial biofilms show resistance against human immune system, as well as against antibiotics. There are great health related concerns because of biofilm potential to cause diseases, utilizing both device-related and non-device-related infections.²

Most hospitalized patients have an intravenous catheter, and up to 30% have a central venous catheter for hemodynamic support, for infusion of medications that require instillation into a central vein, or for long-term venous access.³ These catheters are a major source of health care-associated infections. The Centers for Disease Control and Prevention estimated that 18,000 central lines associated blood stream infections occurred among patients in U.S. intensive care units and 23,000 among patients in inpatient wards in 2009.⁴

Formation of biofilm is universal on central venous catheters, but the location and extent of biofilm formation depend on the duration of catheterization. A short term (less than 10 days) catheters have more biofilm formation on external surface, whereas long term catheters (30 days) have greater biofilm formation on catheter lumen. Growth of microbes may be affected by the nature of the fluid which is administered through central venous catheter.^{5,6} Both Gram negative and Gram positive bacteria can form biofilms on indwelling medical devices. The most common biofilm forming bacteria are *Enterococcus faecalis*, *Staphylococcus aureus*, *S. epidermidis*, *viridans streptococci*, *Escherichia coli*, *Klebsiella Pneumoniae*, *Acinetobacter baumannii* and *Pseudomonas aeruginosa*.⁷

A number of tests are available to detect biofilm production by different bacteria. Methods include Tissue culture plate method, Tube method, Congo red agar, bioluminescent assay and light or fluorescent assay microscopic examination. In this study we screened isolates by tissue culture method to determine their ability to form biofilm.

2. Materials and Methods

This study was carried out from January 2019 to September 2020. During this period, 102 central venous lines were

received in the microbiology laboratory. Positive cultures were obtained from 58 of these 102 samples. Peripheral vein sampling was done to rule out general bacteremia in all these patients to rule out seeding of the catheter through the blood. All cultured catheters were removed using sterile gloves after insertion site has been thoroughly cleaned with 2% povidine –iodine. After catheter was removed, the definition of catheter related infection was based on association of clinical signs (local or systemic) and significant catheter colonization. The distal 5cm of the catheter tip was cut with a sterile surgical scissors and put into separate container and then transported to microbiology laboratory in 2 hours for examination. To improve the detection of biofilm catheter vortexing was done. All catheter tips were cultured by the roll plate culture method directly onto 5% blood agar, Mac Conkey agar and chocolate agar. The plates were incubated at 37°C aerobically for 24 hours. The number of organisms in the plates was evaluated quantitatively. A minimum of 15 colony forming units (CFU) in each plate was considered as positive catheter tip culture. Organisms were identified by standard microbiological procedures including various biochemical tests. All the isolates were subjected to antimicrobial susceptibility tests on Muller-Hinton agar by Kirby Bauer disc diffusion method based on CLSI guidelines.⁸

The isolates were then tested for in vitro production of biofilm using a microtitre plate assay. Biofilm-producing reference strains of *A. baumannii* (ATCC19606) and *P. aeruginosa* (ATCC 27853) were used as positive controls¹⁰ and non-biofilm forming reference strains of *S. aureus* (ATCC25923) and *E. coli* (ATCC25922) as negative controls.

2.1. Biofilm detection method

2.1.1. Tissue culture plate method (TCP)⁹

Organisms isolated from fresh agar plates were inoculated into 10 ml trypticase soy broth with 1% glucose (TSBglu). Broth was incubated at 37°C for 18 hours. The broth culture was then diluted 1:100 with fresh medium. Individual wells of sterile 96 well flat bottom polystyrene tissue culture plates were filled with 200ul of the diluted cultures. Plates were incubated at 37°C for 24 h. contents of each well were removed by gentle tapping. The wells were washed with 0.2 ml of phosphate buffer saline (ph7.2). Biofilm formed by bacteria adherent to the wells was fixed by 2% sodium acetate and stained by crystal violet (0.1%). Excess stain was removed by using deionized water and plates will be kept for drying. Optical density (OD) of stained adherent bacteria was determined with an ELISA reader at wavelength of 570nm. These OD values were considered as an index of bacterial adherence and biofilm formation.

Table 1: Classification of bacterial adherence by TCP method

Mean OD	Adherence	Biofilm Formation
<0.120	Non	Non/weak
0.120-0.240	Moderate	Moderate
>0.240	Strong	High

2.2. Ethical clearance

Ethical clearance was obtained from the Institute's ethical clearance committee.

3. Results

The study was conducted in the clinical bacteriology laboratory, Department of Microbiology of Govt. Medical College, Srinagar and Kashmir. The central venous line received from patients admitted in SICU during January 2019 to December 2020 was processed for isolation and identification of bacterial pathogens according to the standard microbiological techniques.

Demographic characteristics of the study subjects showed that among 102 catheterized patients were isolated (age range 16-65 years) with males 60 (58.82%) and 42(41.1%) females respectively. There was no predilection of infection to any particular gender; patients of both genders were affected, males being marginally more than females. Majority (82%) of the patients were in the age group of 1-60 years, 21-40 yrs (36.27%); 1-20 yrs (27.45%) and 41-60 yrs (18.62%) respectively. A less number of patients were from elderly age groups >61 yrs (17%) as depicted in Table 1.

Table 2: Distribution of cases by age and gender (N= 102)

Age group (years)	Gender		Total No. of cases (n=102)	
	Male	Female	No.	%
1-20	15	13	28	27.45%
21-40	25	12	37	36.27%
41-60	10	9	19	18.62%
61-80	6	5	11	10.78%
>80	4	3	7	6.86%
Total	60 (58.82%)	42 (41.1%)	102	100

Out of 58 isolates, 36 (62.06%) were gram negative bacteria (GNB) and 22(37.93%) were gram positive (Figure 1).

Table 3 depicts that 68% of the patients showed systemic signs of CRBI where as 32% of the patients showed local signs respectively.

Table 4 shows that *Staphylococcus aureus* was most common isolate 16 (27.58%) followed by *Klebsiella* spp. 12 (20.68%), *Acinetobacter* 09 (15.5%), *Coagulase negative Staphylococci* 04 (6.8%) respectively.

Fig. 1: Depicts the bacteriological profile of isolates**Table 3:** Distribution of catheter related bloodstream infection based on clinical signs (N=58)

Local signs	Systemic signs	Total
18(32%)	40(68%)	58(100%)

Table 4: Shows different organisms isolated from central venous line (N=58).

Organisms	No.	%
<i>Staphylococcus aureus</i>	16	27.56
<i>CONS</i>	4	6.89
<i>Enterococcus</i>	1	1.72
<i>Pneumococcus</i>	1	1.72
<i>Acinetobacter</i>	9	15.51
<i>Klebsiella</i>	12	20.68
<i>Pseudomonas</i>	5	8.62
<i>Enterobacter</i>	5	8.62
<i>Citrobacter</i>	2	3.44
<i>E.Coli</i>	3	5.17

Isolates were screened for biofilm formation by TCP method. Out of 58 isolates, classification was done as Strong producer, Moderate producer, Weak/ Non-producer – 20.68%, 27.58%, 51.72% respectively based on their OD values as depicted in Table 5.

Table 5: Screening of the isolates for biofilm formation by tissue culture plate method in gram positive cocci

Organism	Weak	Moderate	Strong	Total
<i>S. aureus</i>	11	2	3	16
<i>CONS</i>	3	1	0	4
<i>Enterococcus</i>	0	1	0	1
<i>Pneumococcus</i>	1	0	0	1

Table 6 depicts screening of the isolates for biofilm formation by tissue culture plate method in gram negative bacilli.

Antibiotic susceptibility patterns of various organisms isolated from central venous line. They showed increased resistance towards penicillin and beta-lactum group of antibiotics, increased sensitivity to linezolid and

Table 6: Shows total number of strong biofilm producers among gram positive cocci

Organism	Biofilm producers(No.)	Percentage
<i>S. aureus</i>	3	18.75
<i>CONS</i>	0	0
<i>Enterococcus</i>	0	0
<i>Pneumococcus</i>	0	0

Table 7: Screening of the isolates for biofilm formation by tissue culture plate method in gram negative bacilli

Organism	Weak	Moderate	Strong	Total
<i>Acinetobacter</i>	5	4	0	9
<i>Pseudomonas</i>	3	0	2	5
<i>Klebsiella</i>	5	3	4	12
<i>E.coli</i>	3	0	0	3
<i>Enterobacter</i>	2	3	0	5
<i>Citrobacter</i>	2	0	0	2

Table 8: Shows total number of strong biofilm producers among gram negative bacilli

Organism	Biofilm producers(No.)	Percentage
<i>Acinetobacter</i>	0	0
<i>Pseudomonas</i>	2	40
<i>Klebsiella</i>	4	33.3
<i>E.coli</i>	0	0
<i>Enterobacter</i>	0	0
<i>Citrobacter</i>	0	0

vancomycin among gram positive organisms. Among gram negative bacteria increased resistance was seen for cephalosporins and aminoglycosides and least resistance for colistin.

Antibiotic sensitivity pattern of different organisms isolated from central venous lines are:

Table 9: Shows antibiotic sensitivity pattern of biofilm producing gram positive bacteria

Organism	Sensitivity pattern
<i>S. aureus</i>	Vancomycin(100%)
	Linezolid(100%)
	Gentamicin(87.5%)
	Cotrimoxazole(75%)
	Clindamycin(62.5%)
	Erythromycin (31.25%)
	Ciprofloxacin (25%)
	Cefoxitin(25%)
	Penicillin (0%)

4. Discussion

Indwelling medical devices are frequently used in all health setting with critical care units of hospitals often using

Table 10: Shows antibiotic sensitivity pattern of biofilmproducing Enterobacteriaceae

Organism	Sensitivity pattern (%)
<i>Klebsiella</i>	Colistin(100%)
	Imipenem(66.6%)
	Ciprofloxacin (66.6%)
	Cotrimoxazole(58.3%)
	Cefepime(50%)
	Gentamicin(50%)
	Piperacillin-Tazobactam(41.6%)
	Ceftazidime(33.3%)
	Tobramycin(33.3%)
<i>E. coli</i>	Colistin(100%)
	Imipenem(100%)
	Amikacin(100%)
	Ciprofloxacin (66%)
	Ceftazidime-Tazobactam(66%)
	Cefepime(33%)
	Tobramycin(33%)
<i>Enterobacter</i>	Colistin (100%)
	Imipenem(100%)
	Amikacin(60%)
	Ticarcillin-Clavulanic-acid (60%)
	Ceftazidime(40%)
	Cefepime(40%)
	Piperacillin-Tazobactam(40%)
	Ciprofloxacin(40%)

Table 11: Shows antibiotic sensitivity pattern of non fermenters

Organism	Sensitivity pattern
<i>Acinetobacter</i>	Colistin(100%)
	Tigecycline(100%)
	Minocycline(66.6%)
	Gentamicin(44.4%)
	Imipenem(33.3%)
	Ciprofloxacin (33.3%)
	Piperacillin-Tazobactam (11.1%)
	Ceftriaxone(11%)
	Cefepime(11%)
<i>Pseudomonas</i>	Colistin(100%)
	Imipenem(80%)
	Ciprofloxacin (80%)
	Gentamicin(60%)
	Piperacillin-Tazobactam(40%)
	Cefepime(40%)
	Ceftazidime-Clavulanic Acid(40%)

multiple medical devices for treatment and intervention in patient care.¹⁰ Intravascular catheters may be inserted for administration of fluids, blood products, medications, nutritional solutions, and hemodynamic monitoring. The organisms may originate from the skin of patients or healthcare workers, to which entry ports are exposed or other sources in the environment.⁷ *Staphylococcus spp.*, *Acinetobacter spp.*, *Klebsiella spp* *Pseudomonas spp.*, and *Enterobacter* are the commonest causes of nosocomial

infections. Colonization of indwelling medical devices with consequent biofilm production is a likely contributory factor to such infections. These microorganisms survive in the hospital environment despite unfavourable conditions such as desiccation, nutrient starvation, and antimicrobial treatment. It is hypothesized that the ability of these microbes to persist in these environments, as well as their virulence may be a result of their capacity to colonize medical devices.¹¹

In the present study, a colonization rate of 53% was recorded in the test catheters. This was consistent with the study of Sadoyama et al (2006),¹² who reported rates of colonization of (23–52%). The predominant contaminants were Gram-negative rods (62.06%). This is in agreement with Liniars et al(1985),¹³ Eyeret al(1990),¹⁴ Greogeret al (1993),¹⁵ Tullu et al(1998),¹⁶ Sherertz et al (2000),¹⁷ Pelletier et al(2000),¹⁸ Hanna et al(2004)¹⁹ and Storti, et al(2005).²⁰

The most prevalent Gram-negative rods isolated were identified as *Klebsiella*(20%), *Acinetobacter*(15%), *Pseudomonas*(8%), *Enterobacter*(8%). Alla et al.²¹ reported that the most prevalent Gram-negative rods isolated were identified as *Acinetobacter baumannii* (23%), followed by *Pseudomonas aeruginosa* (18%), *Klebsiella pneumoniae* (15%) and *Enterobacter* spp. (13%). However, different percentages of other Gram-negative isolates; including *Klebsiella*, *Proteus*, *Pseudomonas*, *Stenotrophomonas*, *Enterobacter*, *Serratia* as well as, *Candida* spp.; were reported by Shapiro et al (1982);²² Flynn et al(1987),²³ Benezra D et al.(1988);²⁴ Sherertz et al(1990),¹⁷ Baleva and Pena, (1997);²⁵ Seifert (1997),²⁶ Tullu et al(1998),¹⁶ Mermel et al (2001),²⁷ Pena et al (2001),²⁸ Cisneros and Rodriguez Bano, (2002),²⁹ Krishnasami et al(2002)³⁰ and Karlowsky et al(2004).³¹ This may be attributed to differences in culture techniques applied, catheterization periods, underlying diseases, types of catheters.

Gram-positive cocci, representing 37.9% of the isolates, were isolated, of these different authors (Darouiche et al 1999³² and Paragioudaki et al 2004³³ reported much higher rates (60–67%).

The total number of positive slime producers, in this study, was 23(39.6%). The highest number of strong slime producer strains was observed in case of *Pseudomonas* (2 out of 5), *Klebsiella pneumoniae* (4 out of 12) and *Staphylococcus* (3 out of 16) Farraget al (2002),³⁴ Banin et al. (2006)³⁵ Sherertz et al (2000)¹⁷ and Walencka et al(2006),³⁶ also reported the ability of *Pseudomonas* for slime production.

The antibiogram of all bacterial isolates was performed using different antibiotics. The results obtained showed that the least resistance rates of Gram-negative bacteria were observed with colistin, tigecycline, imipenem and aminoglycosides. All Gram-positive isolates were sensitive to linezolid, vancomycin followed by aminoglycosides and cotrimoxazoles.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

1. Donlan RM. Biofilm Formation: A Clinically Relevant Microbiological Process. *Clin Infect Dis*. 2001;33(8):1387–92. doi:10.1086/322972.
2. Jamal M, Ahmad W, Andleeb S, Jalil F, Imran M, Nawaz MA, et al. Bacterial biofilm and associated infections. *J Chinese Med Assoc*. 2018;81(1):7–11. doi:10.1016/j.jcma.2017.07.012.
3. Climo M, Diekema D, Warren DK, Herwaldt LA, Perl TM, Peterson L, et al. Prevalence of the Use of Central Venous Access Devices Within and Outside of the Intensive Care Unit: Results of a Survey Among Hospitals in the Prevention Epicenter Program of the Centers for Disease Control and Prevention. *Infect Control Hosp Epidemiol*. 2003;24(12):942–5. doi:10.1086/502163.
4. Vital signs: central line-associated blood stream infections—United States, 2001, 2008, and 2009. *MMWR Morb Mortal Wkly Rep*. 2001;60(8):243–8.
5. Maki DG, Mermel LA. Infections due to infusion therapy. In: Bennett JV, Brachman PS, editors. Hospital infections. Philadelphia: Lippincott-Raven; 1998. p. 689–724.
6. Donlan R, Murga R, Carson L. Growing biofilms in intravenous fluids. In: Wimpenny J, Gilbert P, Walker J, Brading M, editors. Biofilms, the good, the bad, and the ugly. Presented at the fourth meeting of the Biofilm Club. UK: Powys; 1999. p. 23–9.
7. Donlan R. Biofilms and Device-Associated Infections. *Emerg Infect Dis*. 2001;7(2):277–81. doi:10.3201/eid0702.010226.
8. Clinical and Laboratory Standards Institute (CLSI) Performance standards for antimicrobial susceptibility testing; 23rd Informational supplement M100-S23. Wayne: CLSI; 2013.
9. Christensen GD, Simpson WA, Bisno AL, Beachey EH. Adherence of slime-producing strains of *Staphylococcus epidermidis* to smooth surfaces. *Infect Immun*. 1982;37(1):318–26. doi:10.1128/iai.37.1.318-326.1982.
10. Patel FM, Goswami PN, Khara R. Detection of Biofilm formation in device associated clinical bacterial isolates in cancer patients. *Sri Lankan J Infect Dis*. 2016;6(1):43. doi:10.4038/sljid.v6i1.8086.
11. Gelosia A, Baldassarri L, Deighton M, Nguyen T. Phenotypic and genotypic markers of *Staphylococcus epidermidis* virulence. *Clin Microbiol Infect*. 2001;7(4):193–9. doi:10.1046/j.1469-0691.2001.00214.x.
12. Sadoyama G, Filho FG, Filho PP. Central venous catheter-related bloodstream infection caused by *Staphylococcus aureus*: microbiology and risk factors. *Brazilian Journal of Infectious Diseases*. 2006;2(2):100–106.
13. Liñares J, Sitges-Serra A, Garau J, Pérez JL, Martín R. Pathogenesis of catheter sepsis: a prospective study with quantitative and semiquantitative cultures of catheter hub and segments. *J Clin Microbiol*. 1985;21(3):357–60. doi:10.1128/jcm.21.3.357-360.1985.
14. Eyer S, Brummitt C, Crossley K, Siegel R, Cerra F. Catheter related sepsis: prospective randomized trial of three methods of longterm catheter maintenance. *Crit Care Med*. 1990;18:1073–1082.
15. Groeger JS, Lucas AB, Coit D. A prospective, randomised evaluation of the effect of silver impregnated subcutaneous cuffs for preventing tunneled chronic venous access catheter infections in cancer patients. *Am*. 1993;218:206–216.
16. Tullu MS, Deshmukh CT, Baveja SM. Bacterial profile and antimicrobial susceptibility pattern in catheter related nosocomial infection. *Journal of Postgraduate Medicine*. 1998;44(1):7–13.

17. Sherertz RJ. Pathogenesis of vascular catheter infections. In: Waldvogel FA, Bisno A, editors. Infections associated with indwelling medical devices. Washington, D.C: ASM Press; 2000. p. 111–25.
18. Pelletier SJ, Crabtree TD, Gleason TG, Pruett T, Sawyer RG. Bacteremia associated with central venous catheter infection is not an independent predictor of outcomes. No competing interests declared. *J Am Coll Surg*. 2000;190(6):671–80. doi:10.1016/s1072-7515(00)00266-0.
19. Hanna H, Raad I, Darouiche R. New approaches for prevention of intravascular catheter-related infections. *Infect Med*. 2001;18:38–48.
20. Storti A, Pizzolitto AC, Pizzolitto EL. Detection of mixed microbial biofilms on central venous catheters removed from Intensive care Unit Patients. *Braz J Microbiol*. 2005;36(3):275–80. doi:10.1590/s1517-83822005000300013.
21. El-Dien MA, Hosny, Farrag HA, Soheir A, Soheir AA, Hagar S, et al. Prevalence of Central Venous Catheter-Related Infections in Catheterized ICUS' Patients. *Int J Pharm Sci Health Care*. 2014;4(2):67–82.
22. Shapiro ED, Wald ER, Nelson KA, Spiegelman KN. Broviac catheter-related bacteremia in oncology patients. *Am J Dis Child*. 1982;136(8):679–81. doi:10.1001/archpedi.1982.03970440023006.
23. Flynn PM, Shenep JL, Stokes DC. In situ management of confirmed central venous catheter-related bacteremia. *Pediatr Infect Dis J*. 1987;6(8):729–34.
24. Benezra D, Kiehn TE, Gold JWM, Brown AE, Turnbull ADM, Armstrong D. Prospective study of infections in indwelling central venous catheters using quantitative blood cultures. *Am J Med*. 1988;85(4):495–8. doi:10.1016/s0002-9343(88)80084-6.
25. Baleva ME, Pena AC. Catheter-Related Intravascular Infections in Critical Care Units. *Phil J Microbiol Infect Dis*. 1997;26(2):51–4.
26. Seifert H. Catheter-related infections due to gram-negative bacilli. In: Seifert H, Jansen B, Farr BM, editors. Catheter related infections. Marcel Dekker; 1997. p. 111–1138.
27. Mermel LA, Farr BM, Sherertz RJ. Guidelines for the management of intravascular catheter-related infection. *Clin Infect Dis*. 2001;p. 32–1249.
28. Peña C, Pujol M, Ardanuy C, Ricart A, Pallarés R, Liñares J, et al. An outbreak of hospital-acquired *Klebsiella pneumoniae* bacteraemia, including strains producing extended-spectrum β -lactamase. *J Hosp Infect*. 2001;47(1):53–9. doi:10.1053/jhin.2000.0862.
29. Cisneros JM, Rodríguez-Baño J. Nosocomial bacteremia due to *Acinetobacter baumannii*: epidemiology, clinical features and treatment. *Clin Microbiol Infect*. 2002;8(11):687–93. doi:10.1046/j.1469-0691.2002.00487.x.
30. Krishnasami Z, Carlton D, Bimbo L, Taylor ME, Balkovetz DF, Barker J, et al. Management of hemodialysis catheter-related bacteremia with an adjunctive antibiotic lock solution. *Kidney Int*. 2002;61(3):1136–42. doi:10.1046/j.1523-1755.2002.00201.x.
31. Karlowsky JA, Jones ME, Draghi DC, Thornsberry C, Sahm DF, Volturo GA. Prevalence and antimicrobial susceptibilities of bacteria isolated from blood cultures of hospitalized patients in the United States in. *Ann Clin Microbiol Antimicrob*. 2002;3:1–8.
32. Darouiche RO, Raad II, Heard SO, Thornby JJ, Wenker OC, Gabrielli A, et al. A Comparison of Two Antimicrobial-Impregnated Central Venous Catheters. *N Engl J Med*. 1999;340(1):1–8. doi:10.1056/nejm199901073400101.
33. Paragioudaki M, Stamouli V, Kolonitsiou F, Anastassiou ED, Dimitracopoulos G, Spiliopoulou I. Intravenous catheter infections associated with bacteraemia: a 2-year study in a University Hospital. *Clin Microbiol Infect*. 2004;10(5):431–5. doi:10.1111/j.1469-0691.2004.00851.x.
34. Farrag HA, Hussain AA, Salama MS, Mahmoud RR. Molecular analysis, antimicrobial resistance and biological properties contributing to pathogenicity in irradiated clinical isolates associated with urinary tract infection. *Az J Microbiol*. 2002;56:162–81.
35. Banin E, Brady KM, Greenberg EP. Chelator-Induced Dispersal and Killing of *Pseudomonas aeruginosa* Cells in a Biofilm. *Appl Environ Microbiol*. 2006;72(3):2064–9. doi:10.1128/aem.72.3.2064-2069.2006.
36. Walencka E, Sadowska B, Rozalska B. Dual species biofilms: Do *Staphylococci* like other bacteria. *Clin Microbiol Infect*. 2006;12(Suppl. 4):822.

Author biography

Humaira Basher, Lecturer <https://orcid.org/0000-0001-8358-0183>

Anjum Farhana, Professor and Head

Seema Alim, Senior Resident

Shoib Khan, Post Graduate

Cite this article: Basher H, Farhana A, Alim S, Khan S. Detection of catheter related blood stream infections in an ICU of a tertiary care center in Northern India. *Indian J Microbiol Res* 2021;8(2):102-107.